

REC'D 28 APR 2005

WIPO

PA 1260457

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

December 21, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/527,764 ✓

FILING DATE: December 09, 2003 ✓

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



H. L. Jackson

H. L. JACKSON

Certifying Officer

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Himanshu BRAHMBHATT et al.

Title: METHODS FOR TARGETED GENE DELIVERY TO NON-PHAGOCYtic MAMMALIAN CELLS VIA BACTERIALLY DERIVED INTACT MINICELLS *IN-VITRO* AND *IN-VIVO*

Appl. No.: Unassigned

Filing Date: December 9, 2003

15535 U.S. PTO
60/527764
120903

PROVISIONAL PATENT APPLICATION
TRANSMITTAL

Mail Stop PROVISIONAL PATENT APPLICATION
Commissioner for Patents
PO Box 1450
Alexandria, Virginia 22313-1450

Sir:

Transmitted herewith for filing under 37 C.F.R. § 1.53(c) is the provisional patent application of:

Himanshu BRAHMBHATT
Sydney, Australia

Jennifer MACDIARMID
Sydney, Australia

[X] Applicant claims small entity status under 37 CFR 1.27(c)(1).

Enclosed are:

- [X] Specification, Claim(s), and Abstract (44 pages).
- [X] Formal drawings (4 sheets, Figures 1-4).
- [X] Small Entity statement(s).
- [X] Application Data Sheet (37 CFR 1.76).

The filing fee is calculated below:

	Rate	Fee Totals
Basic Fee	\$160.00	\$160.00
[X] Small Entity Fees Apply (subtract ½ of above):	=	\$80.00
TOTAL FILING FEE:	=	\$80.00

[X] A check in the amount of \$80.00 to cover the filing fee is enclosed.

[] The required filing fees are not enclosed but will be submitted in response to the Notice to File Missing Parts of Application.

[X] Further, Applicant, by and through his attorney of record, hereby expressly abandons the application as of the filing date of this letter. This is an abandonment of the application only, and is not to be construed as an abandonment of the invention disclosed in the application. It is respectfully requested that the Office acknowledge abandonment of the application as of the filing date of this letter in a communication mailed to the undersigned.

[X] The Commissioner is hereby authorized to charge any additional fees which may be required regarding this application under 37 C.F.R. §§ 1.16-1.17, or credit any overpayment, to Deposit Account No. 19-0741. Should no proper payment be enclosed herewith, as by a check being in the wrong amount, unsigned, post-dated, otherwise improper or informal or even entirely missing, the Commissioner is authorized to charge the unpaid amount to Deposit Account No. 19-0741.

Please direct all correspondence to the undersigned attorney or agent at the address indicated below.

Respectfully submitted,

Date 9 December 2003 By Stephen A. Bent

FOLEY & LARDNER
Customer Number: 22428
Telephone: (202) 672-5404
Facsimile: (202) 672-5399

Stephen A. Bent
Attorney for Applicant
Registration No. 29,768

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Himanshu BRAHMBHATT et al.

Title: METHODS FOR TARGETED GENE DELIVERY TO NON-
PHAGOCYtic MAMMALIAN CELLS VIA BACTERIALLY
DERIVED INTACT MINICELLS *IN-VITRO* AND *IN-VIVO*

Appl. No.: Unassigned

Filing Date: December 9, 2003

ASSERTION OF SMALL ENTITY STATUS

Commissioner for Patents
PO Box 1450
Alexandria, Virginia 22313-1450

Sir:

Pursuant to 37 C.F.R. §1.27(c)(1), Applicants hereby assert entitlement to small entity status in the above-identified application.

Respectfully submitted,

Date 9 December 2003

By S. A. Bent

FOLEY & LARDNER
Customer Number: 22428
Telephone: (202) 672-5404
Facsimile: (202) 672-5399

Stephen A. Bent
Attorney for Applicant
Registration No. 29,768

U.S. PATENT APPLICATION

for

**METHODS FOR TARGETED GENE DELIVERY TO NON-
PHAGOCYtic MAMMALIAN CELLS VIA BACTERIALLY
DERIVED INTACT MINICELLS *IN-VITRO* AND *IN-VIVO***

Inventors: Himanshu Brahmbhatt
Jennifer MacDiarmid

METHODS FOR TARGETED GENE DELIVERY TO NON-PHAGOCYtic MAMMALIAN CELLS VIA BACTERIALLY DERIVED INTACT MINICELLS

BACKGROUND OF THE INVENTION

5 The present invention relates to a method of targeting bacterial minicell vectors to non-phagocytic host cells, particularly but not exclusively in the context of gene therapy. The method employs bispecific molecules that specifically bind to both a minicell surface structure and a host cell surface structure, such as a receptor. By mediating an interaction between the minicell vectors and non-phagocytic host cells,
10 the bispecific ligands enable targeted delivery of oligonucleotides and polynucleotides to the host cells.

 The objective of gene therapy is to insert one or more foreign genes into the cells of an organism to shut down a gene, to replace a defective gene, or to express a gene product that provides a prophylactic or therapeutic effect. Recent advances in
15 gene therapy have highlighted a variety of methods for introducing foreign genes into the genome of recipient mammals. See Romano *et al.* 1998, 1999; Balicki and Beutler, 2002; Wadhwa *et al.*, 2002; and Thomas *et al.*, 2003. These advances relate to using viral vectors, both human and non-human, and non-viral vectors, such as DNA-liposome complexes.

20 While each vector system has its advantages, each also has significant drawbacks that have limited any clinical application. In particular, viral vectors pose serious safety concerns, including recombination with wild-type viruses, insertional and oncogenic potential, intrinsic toxicity of animal virus vectors to mammalian cells, virus-induced immunosuppression, reversion to virulence of attenuated viruses, and
25 adverse reactions such as an inflammatory response caused by existing immunity. Viral vectors also present practical problems, such as difficulties in recombinant virus manufacture and distribution, low stability, and limited capacity of the vectors to

carry large amounts of exogenous DNA. Non-viral vectors have the drawbacks of generally being less efficient at gene delivery.

Addressing these drawbacks, PCT/IB02/04632 has described recombinant, intact minicells that contain therapeutic nucleic acid molecules. Such minicells are effective vectors for delivering oligonucleotides and polynucleotides to host cells *in vitro* and *in vivo*. PCT/IB02/04632 demonstrated, for example, that recombinant minicells carrying mammalian gene expression plasmids could be delivered to phagocytic cells, such as macrophages, and to non-phagocytic cells, illustrated by human breast cancer cells. The application also showed that intraperitoneal administration of the recombinant minicells resulted in recombinant plasmid delivery to phagocytic cells of the immune system, and that a serum antibody response to the encoded protein could be elicited.

While the efficiency of gene delivery to phagocytic cells via minicells is high (40-60%), the efficiency of gene delivery to non-phagocytic cells heretofore has been comparatively low (3% to 5%). A resultant lack of cell- or organ-specificity has hampered the use of minicell vectors in this regard.

A similar lack of specificity also has hindered the application of non-minicell vectors, and various approaches are under development to address this problem. See Wickham, 2003. One approach makes use of the receptor-mediated endocytosis (RME) system, present in many cell types, and entails development of a diverse set of targeting ligands. In this approach, cell-specificity is imparted on the vector by linking it to a ligand that targets a specific cell surface receptor or marker. Following the specific binding, target cell RME system is activated and the vector/receptor complex is internalized and digested, with some of the DNA payload being transported to the nucleus for gene expression. Some cell receptors may be able to facilitate vector uptake into the cytoplasm directly across the plasma membrane (Fernandez and Bailey, 1998; Phelan *et al.*, 1998; Rojas *et al.*, 1998), but the most common route for receptor-mediated uptake of macromolecular moieties is the endocytic-trafficking pathway (Conner and Schmid, 2003).

Several challenges exist regarding targeted gene delivery to non-phagocytic mammalian cells: (i) breaching the mammalian cell plasma membrane; (ii) exploiting a mechanism of delivery vector internalization; (iii) selecting and understanding the nature of targeting ligands used to target specific mammalian cell surface receptors; 5 (iv) achieving intracellular breakdown of the delivery vector without complete degradation of payload DNA; and (v) obtaining release and transport of payload DNA to the mammalian cell cytoplasm or nucleus. These challenges vary somewhat with each gene delivery vector. Despite intensive research in the field, detailed knowledge of the biological processes involved still is rudimentary.

10 Ligand-based targeting of bacterial cells or any particles of bacterial origin to non-phagocytic cells has not been reported, probably because (a) only live bacterial intracellular pathogens gain entry into non-phagocytic cells by an active invasion process and (b) active cellular invasion would override a passive process such as ligand-based receptor mediated endocytosis. Thus, killed bacterial cells would not 15 engage in active cell invasion, and live bacterial cells would not be directed, contrary to their natural tropism, toward desired non-phagocytic cells. Even if ligand-based targeting was employed to enable endocytosis of killed bacterial cells or non-living particles of bacterial origin, the method would not be expected to be effective for gene delivery. Rather, it would be expected that endosomes would degrade the non-living 20 cells or particles, making them ineffective as gene delivery vectors. In that regard, it currently is thought that only live facultative intracellular pathogenic bacteria can express proteins that allow escape from the endosomal membrane.

To date, no proven methodology exists for effectively targeting bacterial minicell vectors to non-phagocytic mammalian host cells, thereby to deliver a gene 25 payload. Although a variety of vector targeting technologies are known, simply adopting any one of them does not predictably result in a successful, minicell-targeted gene delivery. This is due to the range of biological factors, unique for each gene delivery vector, that can influence targeted gene delivery.

Therefore, a need exists for a method of specifically targeting bacterial minicell vectors to non-phagocytic mammalian cells.

SUMMARY OF THE INVENTION

To address these and other needs, the present invention provides, in accordance with one aspect, a targeted gene delivery method that comprises bringing
5 bispecific ligands into contact with (i) bacterially derived minicells that contain a therapeutic nucleic acid sequence and (ii) non-phagocytic mammalian cells. The ligands have specificity for both a surface component on the minicells and a surface component on the non-phagocytic mammalian cells. As a result, the minicells are
10 engulfed by the mammalian cells, which then produce an expression product of the therapeutic nucleic acid sequence. Contact between the minicells and the mammalian cells may be *in vitro* or *in vivo*.

The invention also provides bispecific ligands useful for targeting minicell vectors to non-phagocytic mammalian host cells. The bispecific ligand may be
15 polypeptide or carbohydrate, and may comprise an antibody or antibody fragment. In preferred embodiments, the bispecific ligand has a first arm that carries specificity for a bacterial minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface structure. A desirable minicell surface structure for ligand binding is an O-polysaccharide component of a
20 lipopolysaccharide. Desirable mammalian cell surface structures for ligand binding are receptors, preferably those capable of activating receptor-mediated endocytosis.

In another aspect, the invention provides a composition comprising (i) bacterially derived minicells that contain a therapeutic nucleic acid and (ii) bispecific
ligands that are capable of binding to a surface component of the minicells and a
25 surface component of a non-phagocytic mammalian cell.

In still another aspect, the invention provides for the use of bacterially derived minicells that contain a therapeutic nucleic acid and a bispecific ligand in the preparation of a medicament for use in a method of treating a disease or modifying a

trait by administration of the medicament to a cell, tissue, or organ. Such medicaments are useful to treat various conditions and diseases by increasing expression or function of a desired protein, or by inhibiting expression or function of target protein. The disease to be treated in this context may be a cancer, for example, or an acquired disease, such as AIDS, pneumonia, emphysema, or a condition engendered by an inborn error of metabolism, such as cystic fibrosis. Alternatively, the treatment may affect a trait, such as fertility, or an immune response associated with an allergen or an infectious agent.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows efficient internalization of human androgen receptor-targeted recombinant minicells, in contrast to non-targeted minicells, into human prostate carcinoma LNCaP cells. The procedures were performed as described in Example 1 and the results were visualized by confocal microscopy. Immunofluorescence staining was performed for all shown images with anti-*S. typhimurium* LPS specific monoclonal antibody, followed by Alexa Fluor 594-conjugated goat anti-mouse IgG (H+L) antibody. Each figure is shown as an overlap of Differential Interference Contrast (DIC) and red fluorescence images. (A) Control LNCaP cells not transfected with minicells. No red fluorescence was observed following staining for *S. typhimurium* LPS. (B) LNCaP cells transfected with non-targeted minicells and stained after 16 hr co-incubation. Very few background red fluorescence dots were observed. (C) LNCaP cells transfected with targeted minicells and stained after 16hrs. Most cells showed red fluorescence in the cytoplasm, revealed in the black-and-white image as light grey. (D) LNCaP cells transfected with non-targeted minicells and stained after 24 hr co-incubation. Very few background red fluorescence dots were observed. (E) LNCaP cells transfected with targeted minicells and stained after 24 hrs. The result showed intense red fluorescence in the cytoplasm of most cells (light grey in the image), (F) Same as (E) but at a higher magnification to show a single transfected cell. Almost all the cytoplasm fluoresced red (light grey). Scale bars are shown for each image.

Figure 2 shows efficient internalization of EGF receptor-targeted recombinant minicells, *versus* non-targeted minicells, into human breast cancer MDA-MB-468 cells. The procedures were performed as described in Example 2 and the results were visualized by confocal microscopy. Immunofluorescence staining was performed for all shown images with anti-*S. typhimurium* LPS specific monoclonal antibody, followed by Alexa Fluor 594-conjugated goat anti-mouse IgG (H+L) antibody. Each image is shown as an overlap of DIC and red fluorescence images. (A) Control MDA-MB-468 cells not transfected with minicells. No red fluorescence was observed following staining for *S. typhimurium* LPS. (B) MDA-MB-468 cells transfected with non-targeted minicells and stained after 24 hr co-incubation. Very few background red fluorescence dots were observed. (C) MDA-MB-468 cells transfected with targeted minicells and stained after 24 hrs. Most cells showed red fluorescence on the surface and some in the cytoplasm (light grey area in the black-white image). (D) Same as (C) but at a higher magnification to reveal a single cell. The result was the same as for (C). (E) Same as (D) except cells were stained after 36 hrs. The result showed intense red fluorescence in the cytoplasm of most cells (light grey in the image). Scale bars are shown for each image.

Figure 3 shows efficient internalization of Her2/neu receptor-targeted recombinant minicells, *versus* non-targeted minicells, into human ovarian cancer SKOV-3 cells. The procedures were performed as described in Example 3 and the results were visualized by confocal microscopy. Immunofluorescence staining was performed for all shown images with anti-*S. typhimurium* LPS specific monoclonal antibody, followed by Alexa Fluor 594-conjugated goat anti-mouse IgG (H+L) antibody. Each image is shown as an overlap of DIC and red fluorescence images. (A) Control SKOV-3 cells not transfected with minicells. No red fluorescence was observed following staining for *S. typhimurium* LPS. (B) SKOV-3 cells transfected with non-targeted minicells and stained after 36 hr co-incubation. Very few background red fluorescence dots were observed. (C) SKOV-3 cells transfected with targeted minicells and stained after 36 hrs. Most cells showed red fluorescence in the cytoplasm (light grey area in the black-white image). (D) Same as (C) but at a higher magnification. The result was the same as for (C). (E) Same as (C) but higher

magnification to show a few cells. The result showed intense red fluorescence in the cytoplasm of most cells (light grey in the image). Scale bars are shown for each image.

Figure 4 shows treatment of human breast cancer xenografts in nude mice via
 5 targeted recombinant minicells. Breast cancer xenografts were established in nude
 mice (see example 4) and treated intratumorally with targeted recombinant minicells
 carrying plasmid pORF5-HSV1tk::Sh ble. **(Group 1, control)** tumors did not receive
 any treatment; **(Group 2, control)** tumors were treated with non-targeted recombinant
 minicells [M-HSVtk] followed by 2 doses of GCV; **(Group 3, control)** tumors were
 10 treated with targeted recombinant minicells [TM-HSVtk]; **(Group 4, control)** tumors
 were treated with the bispecific antibody (BsAb; anti-*S. typhimurium* LPS / anti-
 human EGF receptor specificities), followed by 2 doses of GCV; **(Group 5,**
experimental) tumors were treated with targeted recombinant minicells [TM-HSVtk]
 followed by one dose of GCV; **(Group 6, experimental)** tumors were treated with
 15 targeted recombinant minicells [TM-HSVtk] followed by 2 doses of GCV. Below the
 x-axis are shown the days on which various treatments were given to specific groups.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present inventors have discovered that bispecific ligands can be employed
 to target bacterial minicell vectors to non-phagocytic mammalian host cells. Such
 20 host cells normally are resistant to adhesion and endocytosis of minicells *in-vivo*, yet
 can be made receptive to minicell delivery vector binding and internalization with the
 aid of a bispecific ligand.

Additionally, the inventors have discovered that the internalized minicells
 were degraded sufficiently to release recombinant plasmid DNA. This was surprising
 25 because non-phagocytic mammalian cells inherently do not carry aggressive
 intracellular compartments like phagolysosomes, which predominantly exist in cells
 of the immune system such as phagocytic macrophages.

As an additional surprise, the inventors also discovered that bacterial minicells can effect recombinant plasmid escape from the late-endosome of non-phagocytic cells. This was unexpected because minicells are non-living and devoid of the parent bacterial chromosome that encodes late-endosomal and phagosomal membrane-lysing proteins. Indeed, it had been commonly accepted that only live facultative intracellular bacterial pathogens designed to lyse the lysosomal membrane and release DNA intracellularly can deliver genes to non-professional phagocytes (reviewed recently by Grillot-Courvalin *et al.*, 2002). For example, *Listeria monocytogenes* expresses a pore-forming cytolysin, Listeriolysin O (chromosomally encoded by the *hly* gene), that is thought to play a major role in lysing the endosomal and phagosomal membrane, thereby allowing recombinant DNA to enter an infected cell cytoplasm. Similarly, *Shigella flexneri* is also thought to escape the phagocytic vacuole by lysing the phagosomal membrane.

The inventors further have established that effective minicell-mediated recombinant gene delivery to the nucleus of non-phagocytic cells relates to the number of plasmid copies carried by a minicell. Thus, minicells carrying a high-copy number plasmids (over 60 plasmid copies per minicell) effect efficient gene delivery to non-phagocytic cells, whereas minicells carrying a medium-copy (11 to 60 per minicell) or low-copy (1 to 10 per minicell) number plasmids are less effective.

Additionally, the inventors have established that efficiency of gene delivery relates to the number of recombinant minicells that are endocytosed within endosomes. Accordingly, non-phagocytic target cells that carry highly expressed receptors on the cell surface, such as EGF receptor on the surface of some human breast cancer cells and to which the bispecific ligand was targeted, show more minicells engulfed within each endosome, often more than 10, resulting in highly efficient recombinant gene delivery to the cell nucleus. These results suggest that the chances for escape of recombinant DNA from late endosomes are increased when the recombinant DNA load within an endosome is high enough to compensate for losses through degradation within the endosome. The results also show that effective gene delivery may be achieved by exploiting mammalian cell surface receptors that are

over-expressed on the cell surface, thereby enabling the endocytosis of multiple minicells within individual endosomes.

In accordance with the foregoing discoveries, the invention broadens the spectrum of diseases amenable to gene therapy using minicell vectors, by enhancing
5 the minicell transfection efficiency in target cells or tissues that normally are refractory to minicell adhesion, endocytosis and gene delivery. The ability to target minicells also provides a safer and more flexible system for gene therapy.

In one aspect, therefore, the invention provides a targeted gene delivery method that comprises bringing bispecific ligands into contact with (a) bacterially
10 derived minicells that contain a therapeutic nucleic acid sequence and (b) non-phagocytic mammalian cells. The bispecific ligands, having specificity for both minicell and mammalian cell components, cause the minicells to bind to the mammalian cells, such that the minicells are engulfed by the mammalian cells, which then produce an expression product of the therapeutic nucleic acid sequence.

The inventors found that this method is broadly applicable to a range of non-
15 phagocytic mammalian cells that normally are refractory to specific adhesion and endocytosis of minicells. For example, bispecific antibody ligands with anti-O-polysaccharide specificity on one arm and anti-HER2 receptor, anti-EGF receptor or anti-androgen receptor specificity on the other arm efficiently bound minicells to the
20 respective receptors on a range of non-phagocytic cells. These cells included lung, ovarian, brain, breast, prostate and skin cancer cells. Moreover, the efficient binding preceded rapid endocytosis of the minicells by each of the non-phagocytic cells.

The inventors' discovery was surprising because it previously was thought that only "professional" phagocytes, such as macrophages and neutrophils, can endocytose
25 large macromolecular particles like bacterial cells, which are 600 nm and larger. Conversely, it was thought that non-phagocytic mammalian cells can endocytose only small, non-living macromolecular particles such as liposomes, which are 150-400 nm, and viruses, which are on the order of 65-80 nm in size. See Bondoc and Fitzpatrick,

1998. Bacterially derived intact minicells used in the inventors' studies were approximately 500 nm in diameter.

The inventors also found that recombinant DNA carried by minicells can be expressed by non-phagocytic mammalian host cells. The minicells, once
5 endocytosed, subsequently become degraded in late endosomes. Some recombinant DNA carried by the minicells, however, escapes the endosomal membranes and is transported to the mammalian cell nucleus, permitting gene expression. This discovery was surprising because it previously was thought that only live facultative intracellular pathogens carry virulence proteins capable of endosomal membrane
10 escape and gene delivery. See Grillot-Courvalin *et al.*, 2002. Non-living bacteria or bacterially derived minicells were not expected to express these *in-vivo* induced virulence proteins and, hence, were expected to be completely degraded within endosomes, with no possibility for endosomal escape by any recombinant DNA.

The invention therefore provides novel methods that extend the range of
15 mammalian cells amenable to gene therapy via bacterially derived minicells. These methods may be performed both *in vitro* and *in vivo*.

Ligands useful in the invention include any agent that binds to a surface component on a target cell and to a surface component on a minicell. Preferably, the surface component on a target cell is a receptor, especially a receptor capable of
20 mediating endocytosis. The ligands may comprise a polypeptide and/or carbohydrate component. Antibodies are preferred ligands. For example, a bispecific antibody that carries dual specificities for a surface component on bacterially derived intact minicells and for a surface component on target mammalian cells, can be used to efficiently target the minicells to the target mammalian cells *in vitro* and *in vivo*.
25 Useful ligands also include receptors, enzymes, binding peptides, fusion/chimeric proteins and small molecules.

The selection of a particular ligand is made on two primary bases: (i) specific binding to one or more domains on the surface of intact minicells and (ii) specific binding to one or more domains on the surface of the target cells. Thus, ligands

preferably have a first arm that carries specificity for a bacterially derived intact minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface structure. Each of the first and second arms may be multivalent. Preferably, each arm is monospecific, even if multivalent.

5 For binding to bacterially derived minicells, it is desirable for one arm of the ligand to be specific for the O-polysaccharide component of a lipopolysaccharide found on the parent bacterial cell. Other minicell surface structures that can be exploited for ligand binding include cell surface-exposed polypeptides and carbohydrates on outer membranes, pilli, fimbriae and flagella.

10 For binding to target cells, one arm of the ligand is specific for a surface component of a non-phagocytic mammalian cell. Such components include cell surface proteins, peptides and carbohydrates, whether characterized or uncharacterized. Cell surface receptors, especially those capable of activating receptor-mediated endocytosis, are desirable cell surface components for targeting.

15 By way of example, one may target tumor cells, metastatic cells, vasculature cells, such as endothelial cells and smooth muscle cells, lung cells, kidney cells, blood cells, bone marrow cells, brain cells, liver cells, and so forth, or precursors of any selected cell by selecting a ligand that specifically binds a cell surface receptor motif on the desired cells. Examples of cell surface receptors include carcinoembryonic
20 antigen (CEA), which is overexpressed in most colon, rectum, breast, lung, pancreas and gastrointestinal tract carcinomas (Marshall, 2003); heregulin receptors (HER-2, *neu* or *c-erbB-2*), which is frequently overexpressed in breast, ovarian, colon, lung, prostate and cervical cancers (Hung *et al.*, 2000); epidermal growth factor receptor (EGFR), which is highly expressed in a range of solid tumors including those of the
25 breast, head and neck, non-small cell lung and prostate (Salomon *et al.*, 1995); asialoglycoprotein receptor (Stockert, 1995); transferrin receptor (Singh, 1999); serpin enzyme complex receptor, which is expressed on hepatocytes (Ziady *et al.*, 1997); fibroblast growth factor receptor (FGFR), which is overexpressed on pancreatic ductal adenocarcinoma cells (Kleeff *et al.*, 2002); vascular endothelial growth factor receptor

(VEGFR), for anti-angiogenesis gene therapy (Becker *et al.*, 2002 and Hoshida *et al.*, 2002); folate receptor, which is selectively overexpressed in 90% of nonmucinous ovarian carcinomas (Gosselin and Lee, 2002); cell surface glycocalyx (Batra *et al.*, 1994); carbohydrate receptors (Thurnher *et al.*, 1994); and polymeric immunoglobulin receptor, which is useful for gene delivery to respiratory epithelial cells and attractive for treatment of lung diseases such as Cystic Fibrosis (Kaetzel *et al.*, 1997).

Preferred ligands comprise antibodies and/or antibody derivatives. As used herein, the term "antibody" encompasses an immunoglobulin molecule obtained by *in vitro* or *in vivo* generation of an immunogenic response. The term "antibody" includes polyclonal, monospecific and monoclonal antibodies, as well as antibody derivatives, such as single-chain antibody fragments (scFv). Antibodies and antibody derivatives useful in the present invention also may be obtained by recombinant DNA techniques.

Wild type antibodies have four polypeptide chains, two identical heavy chains and two identical light chains. Both types of polypeptide chains have constant regions, which do not vary or vary minimally among antibodies of the same class, and variable regions. Variable regions are unique to a particular antibody and comprise an antigen binding domain that recognizes a specific epitope. The regions of the antigen binding domain that are most directly involved in antibody binding are "complementarity-determining regions" (CDRs).

The term "antibody" also encompasses derivatives of antibodies, such as antibody fragments that retain the ability to specifically bind to antigens. Such antibody fragments include Fab fragments (a fragment that contains the antigen-binding domain and comprises a light chain and part of a heavy chain bridged by a disulfide bond), Fab' (an antibody fragment containing a single antigen-binding domain comprising a Fab and an additional portion of the heavy chain through the hinge region, F(ab')₂ (two Fab' molecules joined by interchain disulfide bonds in the hinge regions of the heavy chains), a bispecific Fab (a Fab molecule having two antigen binding domains, each of which may be directed to a different epitope), and

an scFv (the variable, antigen-binding determinative region of a single light and heavy chain of an antibody linked together by a chain of amino acids.)

When antibodies, including antibody fragments, constitute part or all of the ligands, they preferably are of human origin or are modified to be suitable for use in humans. So-called "humanized antibodies" are well known in the art. See, *e.g.*, Osbourn *et al.*, 2003. They have been modified by genetic manipulation and/or *in vitro* treatment to reduce their antigenicity in a human. Methods for humanizing antibodies are described, *e.g.*, in U.S. patents No. 6,639,055, No. 5,585,089, and No. 5,530,101. In the simplest case, humanized antibodies are formed by grafting the antigen-binding loops, known as complementarity-determining regions (CDRs), from a mouse mAb into a human IgG. See Jones *et al.*, 1986; Riechmann *et al.*, 1988; and Verhoeven *et al.*, 1988. The generation of high-affinity humanized antibodies, however, generally requires the transfer of one or more additional residues from the so-called framework regions (FRs) of the mouse parent mAb. Several variants of the humanization technology also have been developed. See Vaughan *et al.*, 1998.

Human antibodies, rather than "humanized antibodies," also may be employed in the invention. They have high affinity for their respective antigens and are routinely obtained from very large, single-chain variable fragments (scFvs) or Fab phage display libraries. See Griffiths *et al.*, 1994; Vaughan *et al.*, 1996; Sheets *et al.*, 1998; de Haard *et al.*, 1999; and Knappik *et al.*, 2000.

Useful ligands also include bispecific single chain antibodies, which typically are recombinant polypeptides consisting of a variable light chain portion covalently attached through a linker molecule to a corresponding variable heavy chain portion. See U.S. Patents 5,455,030; 5,260,203 and 4,496,778. Bispecific antibodies also can be made by other methods. For example, chemical heteroconjugates can be created by chemically linking intact antibodies or antibody fragments of different specificities. See Karpovsky *et al.*, 1984. However, such heteroconjugates are difficult to make in a reproducible manner and are at least twice as large as normal monoclonal antibodies. Bispecific antibodies also can be created by disulfide

exchange, which involves enzymatic cleavage and reassociation of the antibody fragments. See Glennie *et al.*, 1987.

Because Fab and scFv fragments are monovalent they often have low affinity for target structures. Therefore, preferred ligands made from these components are engineered into dimeric, trimeric or tetrameric conjugates to increase functional affinity. See Tomlinson and Holliger, 2000; Carter, 2001; Hudson and Souriau, 2001; and Todorovska *et al.*, 2001. Such conjugate structures may be created by chemical and/or genetic cross-links.

Bispecific ligands of the invention preferably are monospecific at each end, *i.e.*, specific for a single component on minicells at one end and specific for a single component on target cells at the other end. The ligands may be multivalent at one or both ends, for example, in the form of so-called diabodies, triabodies and tetrabodies. See Hudson and Souriau, 2003. A diabody is a bivalent dimer formed by a non-covalent association of two scFvs, which yields two Fv binding sites. Likewise, a triabody results from the formation of a trivalent trimer of three scFvs, yielding three binding sites, and a tetrabody results from the formation of a tetravalent tetramer of four scFvs, yielding four binding sites.

Several humanized, human, and mouse monoclonal antibodies and fragments thereof that have specificity for receptors on mammalian cells have been approved for human therapeutic use, and the list is growing rapidly. See Hudson and Souriau, 2003. An example of such an antibody that can be used to form one arm of a bispecific ligand has specificity for HER2: HerceptinTM; Trastuzumab.

Antibody variable regions also can be fused to a broad range of protein domains. Fusion to human immunoglobulin domains such as IgG1 CH3 both adds mass and promotes dimerization. See Hu *et al.*, 1996. Fusion to human Ig hinge-Fc regions can add effector functions. Also, fusion to heterologous protein domains from multimeric proteins promotes multimerization. For example, fusion of a short scFv to short amphipathic helices has been used to produce miniantibodies. See Pack and Pluckthun, 1992. Domains from proteins that form heterodimers, such as fos/jun, can

be used to produce bispecific molecules (Kostelny *et al.*, 1992) and, alternately, homodimerization domains can be engineered to form heterodimers by engineering strategies such as “knobs into holes” (Ridgway *et al.*, 1996). Finally, fusion protein partners can be selected that provide both multimerization as well as an additional function, *e.g.* streptavidin. See Dubel *et al.*, 1995.

Minicells of the invention are anucleate forms of *E. coli* or other bacterial cells, engendered by a disturbance in the coordination, during binary fission, of cell division with DNA segregation. Prokaryotic chromosomal replication is linked to normal binary fission, which involves mid-cell septum formation. In *E. coli*, for example, mutation of *min* genes, such as *minCD*, can remove the inhibition of septum formation at the cell poles during cell division, resulting in production of a normal daughter cell and an anucleate minicell. See de Boer *et al.*, 1992; Raskin & de Boer, 1999; Hu & Lutkenhaus, 1999; Harry, 2001. Minicells are distinct from other small vesicles that are generated and released spontaneously in certain situations and, in contrast to minicells, are not due to specific genetic rearrangements or episomal gene expression. For practicing the present invention, it is desirable for minicells to have intact cell walls (“intact minicells”).

In addition to *min* operon mutations, anucleate minicells also are generated following a range of other genetic rearrangements or mutations that affect septum formation, for example in the *divIVB1* in *B. subtilis*. See Reeve and Cornett, 1975; Levin *et al.*, 1992. Minicells also can be formed following a perturbation in the levels of gene expression of proteins involved in cell division/chromosome segregation. For example, overexpression of *minE* leads to polar division and production of minicells. Similarly, chromosome-less minicells may result from defects in chromosome segregation for example the *smc* mutation in *Bacillus subtilis* (Britton *et al.*, 1998), *spoOJ* deletion in *B. subtilis* (Ireton *et al.*, 1994), *mukB* mutation in *E. coli* (Hiraga *et al.*, 1989), and *parC* mutation in *E. coli* (Stewart and D’Ari, 1992). Gene products may be supplied *in trans*. When over-expressed from a high-copy number plasmid, for example, CafA may enhance the rate of cell division and/or inhibit chromosome partitioning after replication (Okada *et al.*, 1994), resulting in formation of chained

cells and anucleate minicells (Wachi *et al.*, 1989; Okada *et al.*, 1993). Minicells can be prepared from any bacterial cell of Gram-positive or Gram-negative origin.

Minicells of the invention contain a nucleic acid molecule that, upon transcription and/or translation, function to ameliorate or otherwise treat a disease or modify a trait in a cell, tissue or organ. For purposes of the present description, such nucleic acid molecules are categorized as "therapeutic nucleic acid molecules." Ordinarily, the therapeutic nucleic acid is found on a plasmid within the minicells.

The therapeutic nucleic acid molecule encodes a product, such as functional RNA (*e.g.*, antisense or siRNA) or a peptide, polypeptide or protein, the production of which is desired. For example, the genetic material of interest can encode a hormone, receptor, enzyme, or (poly) peptide of therapeutic value. Such methods can result in transient expression of non-integrated transferred DNA, extrachromosomal replication and expression of transferred replicons such as episomes, or integration of transferred genetic material into the genomic DNA of host cells.

Transcription or translation of a given therapeutic nucleic acid molecule may be useful in treating cancer or an acquired disease, such as AIDS, pneumonia, emphysema, or in correcting inborn errors of metabolism, such as cystic fibrosis. Transcription or translation of a therapeutic nucleic acid may also effect contraceptive sterilization, including contraceptive sterilization of feral animals. Allergen-mediated and infectious agent-mediated inflammatory disorders also can be countered by administering, via the present invention, a therapeutic nucleic acid molecule that, upon expression in a patient, affects immune response(s) associated with the allergen and infectious agent, respectively. A therapeutic nucleic acid molecule also may have an expression product, or there may be a downstream product of post-translational modification of the expression product, that reduces the immunologic sequelae related to transplantation or that helps facilitate tissue growth and regeneration.

A therapeutic nucleic acid molecule may be the normal counterpart of a gene that expresses a protein that functions abnormally or that is present in abnormal levels in a disease state, as is the case, for example, with the cystic fibrosis transmembrane

conductance regulator in cystic fibrosis (Kerem *et al.*, 1989; Riordan *et al.*, 1989; Rommens *et al.*, 1989), with β -globin in sickle-cell anemia, and with any of α -globin, β -globin and γ -globin in thalassemia. The therapeutic nucleic acid molecule can have an antisense RNA transcript or small interfering RNA, as mentioned above. Thus, an
5 excess production of α -globin over β -globin which characterizes β -thalassemia can be ameliorated by gene therapy, in accordance with the present invention, using an intact minicell engineered to contain a plasmid incorporating a sequence that has an antisense RNA transcript vis-à-vis a target sequence of the α -globin mRNA.

In the treatment of cancer, a therapeutic nucleic acid molecule suitable for use
10 according to the present invention could have a sequence that corresponds to or is derived from a gene that is associated with tumor suppression, such as the *p53* gene, the retinoblastoma gene, and the gene encoding tumor necrosis factor. A wide variety of solid tumors -- cancer, papillomas, and warts -- should be treatable by this approach, pursuant to the invention. Representative cancers in this regard include
15 colon carcinoma, prostate cancer, breast cancer, lung cancer, skin cancer, liver cancer, bone cancer, ovary cancer, pancreas cancer, brain cancer, head and neck cancer, and lymphoma. Illustrative papillomas are squamous cell papilloma, choroid plexus papilloma and laryngeal papilloma. Examples of wart conditions are genital warts, plantar warts, epidermodysplasia verruciformis, and malignant warts.

20 A therapeutic nucleic acid molecule for the present invention also can comprise a DNA segment coding for an enzyme that converts an inactive prodrug into one or more cytotoxic metabolites so that, upon *in vivo* introduction of the prodrug, the target cell in effect is compelled, perhaps with neighboring cells as well, to commit suicide. Preclinical and clinical applications of such a "suicide gene," which
25 can be of non-human origin or human origin, are reviewed by Spencer (2000), Shangara *et al.* (2000) and Yazawa *et al.* (2002). Illustrative of suicide genes of non-human origin are those that code for HSV-thymidine kinase (*tk*), cytosine deaminase (CDA) + uracil phosphoribosyltransferase, xanthine-guanine phosphoribosyl-transferase (GPT), nitroreductase (NTR), purine nucleoside phosphorylase (PNP, DeoD),
30 cytochrome P450 (CYP4B1), carboxypeptidase G2 (CPG2), and D-amino acid

oxidase (DAAO), respectively. Human-origin suicide genes are exemplified by genes that encode carboxypeptidase A1 (CPA), deoxycytidine kinase (dCK), cytochrome P450 (CYP2B1,6), LNGFR/FKBP/Fas, FKBP/Caspases, and ER/p53, respectively.

A suicide-gene therapy could be applied to the treatment of AIDS. This strategy has been tested with suicide vectors that express a toxic gene product as soon as treated mammalian cells become infected by HIV-1. These vectors use the HIV-1 regulatory elements, Tat and/or Rev, to induce the expression of a toxic gene such as α -diphtheria toxin, cytosine deaminase, or interferon- α 2 after infection by HIV-1. See Curiel *et al.*, 1993; Dinges *et al.*, 1995; Harrison *et al.*, 1992a; Harrison *et al.*, 1992b; Ragheb *et al.*, 1999.

The therapeutic nucleic acid of the invention typically is contained on a plasmid within the minicell. The plasmid also may contain an additional nucleic acid segment that functions as a regulatory element, such as a promoter, a terminator, an enhancer or a signal sequence, and that is operably linked to the therapeutic nucleic acid segment. A suitable promoter can be tissue-specific or even tumor-specific, as the therapeutic context dictates.

A promoter is "tissue-specific" when it is activated preferentially in a given tissue and, hence, is effective in driving expression, in the target tissue, of an operably linked structural sequence. The category of tissue-specific promoters includes, for example: the hepatocyte-specific promoter for albumin and α_1 -antitrypsin, respectively; the elastase I gene control region, which is active in pancreatic acinar cells; the insulin gene control region, active in pancreatic beta cells; the mouse mammary tumor virus control region, which is active in testicular, breast, lymphoid and mast cells; the myelin basic protein gene control region, active in oligodendrocyte cells in the brain; and the gonadotropic releasing hormone gene control region, which is active in cells of the hypothalamus. See Frain *et al.* (1990), Ciliberto *et al.* (1985), Pinkert *et al.*, (1987), Kelsey *et al.* (1987), Swift *et al.* (1984), MacDonald (1987), Hanahan, (1985), Leder *et al.* (1986), Readhead *et al.* (1987), and Mason *et al.* (1986).

There also are promoters that are expressed preferentially in certain tumor cells or in tumor cells *per se*, and that are useful for treating different cancers in accordance with the present invention. The class of promoters that are specific for cancer cells is illustrated by: the tyrosinase promoter, to target melanomas; the MUC1/Df3 promoter, to target breast carcinoma; the hybrid *myoD* enhancer/SV40 promoter, which targets expression to rhabdomyosarcoma (RMS); the carcinoembryonic antigen (CEA) promoter, which is specific for CEA-expressing cells such as colon cancer cells, and the hexokinase type II gene promoter, to target non-small cell lung carcinomas. See Hart (1996), Morton & Potter (1998), Kurane *et al.* (1998) and Katabi *et al.* (1999).

A signal sequence can be used, according to the present invention, to effect secretion of an expression product or localization of an expression product to a particular cellular compartment. Thus, a therapeutic polynucleotide molecule that is delivered via intact minicells may include a signal sequence, in proper reading frame, such that the expression product of interest is secreted by an engulfing cell or its progeny, thereby to influence surrounding cells, in keeping with the chosen treatment paradigm. Illustrative signal sequences include the haemolysin C-terminal secretion sequence, described in U.S. patent No. 5,143,830, the BAR1 secretion sequence, disclosed in U.S. patent No. 5,037,743, and the signal sequence portion of the zsig32 polypeptide, described in U.S. patent No. 6,025,197.

A plasmid within a minicell of the invention also may contain a reporter element. A reporter element confers on its recombinant host a readily detectable phenotype or characteristic, typically by encoding a polypeptide, not otherwise produced by the host, that can be detected, upon expression, by histological or *in situ* analysis, such as by *in vivo* imaging techniques. For example, a reporter element delivered by an intact minicell, according to the present invention, could code for a protein that produces, in the engulfing host cell, a colorimetric or fluorometric change that is detectable by *in situ* analysis and that is a quantitative or semi-quantitative function of transcriptional activation. Illustrative of these proteins are esterases,

phosphatases, proteases and other enzymes, the activity of which generates a detectable chromophore or fluorophore.

Preferred examples are *E. coli* β -galactosidase, which effects a color change via cleavage of an indigogenic substrate, indolyl- β -D-galactoside, and a luciferase, which oxidizes a long-chain aldehyde (bacterial luciferase) or a heterocyclic carboxylic acid (luciferin), with the concomitant release of light. Also useful in this context is a reporter element that encodes the green fluorescent protein (GFP) of the jellyfish, *Aequorea victoria*, as described by Prasher *et al.* (1995). The field of GFP-related technology is illustrated by two published PCT applications, WO 095/21191 (discloses a polynucleotide sequence encoding a 238 amino-acid GFP apoprotein, containing a chromophore formed from amino acids 65 through 67) and WO 095/21191 (discloses a modification of the cDNA for the apo-peptide of *A. victoria* GFP, providing a peptide having altered fluorescent properties), and by a report of Heim *et al.* (1994) of a mutant GFP, characterized by a 4-to-6-fold improvement in excitation amplitude.

Another type of a reporter element is associated with an expression product that renders the recombinant minicell resistant to a toxin. For instance, the *neo* gene protects a host against toxic levels of the antibiotic G418, while a gene encoding dihydrofolate reductase confers resistance to methotrexate, and the chloramphenicol acetyltransferase (CAT) gene bestows resistance to chloramphenicol.

Other genes for use as a reporter element include those that can transform a host minicell to express distinguishing cell-surface antigens, *e.g.*, viral envelope proteins such as HIV gp120 or herpes gD, which are readily detectable by immunoassays.

Target cells of the invention include any cell into which an exogenous nucleic acid molecule is to be introduced. ("Introduced," when used in reference to an exogenous nucleic acid molecule, means that the nucleic acid molecule carried within a minicell is delivered to the target cell.) Desirable target cells are characterized by expression of a cell surface receptor that, upon binding of a ligand, facilitates

endocytosis. Preferred target cells are non-phagocytic, meaning that the cells ordinarily do not ingest bacterial particles, and are mammalian.

Methods of the invention can be used to deliver a range of nucleic acid molecules, which can be cDNA as well as genomic DNA or RNA, and can be in the sense or the anti-sense orientation. The nucleic acid molecule present in a minicell, pursuant to the present invention, can take the form of a plasmid, expression vector, or other genetic construct, but is not genomic DNA originating from the bacterial cell that gave rise to the minicell. Suitable for use in the present invention is any desired DNA or RNA sequence from a eukaryotic, prokaryotic, or synthetic source which may be placed under the translational and transcriptional control of a eukaryotic gene expression promoter, or which may be expressed in the mammalian cell using trans-activating factors from the host cell.

Methods of the invention may be performed *in vivo* or *ex vivo*. In an *ex vivo* procedure, for example, target cells may be removed from a subject, such as by biopsy. An appropriate ligand may be selected based on knowledge of a cell surface receptor that is expressed by the target cells. The gene(s) to be delivered to the target cells are cloned into an appropriate episomal carrier DNA, for example a plasmid, and transferred into parent bacterial cells from which the intact minicells are to be derived. Processes for obtaining minicells are well known in the art, as described in PCT/IB02/04632. Minicells carrying the recombinant DNA are then purified by procedures known in the art and described in PCT/IB02/04632. The bispecific ligand is then bound to the recombinant purified minicells, for example by *in-vitro* incubation in suitable medium, and excess ligand is washed away from the ligand-loaded minicells. The composition comprising purified intact minicells and the bispecific ligand, attached to the minicells via one arm that has specificity for a minicell surface component, is then brought into contact with target cells either *in vitro*, for example in tissue culture (as described in Example 1, 2 and 3), or *in vivo* (as described in example 4).

Thus, the invention includes a method for performing *ex vivo* gene therapy into desired non-phagocytic mammalian cells that are normally refractory to minicell-mediated gene therapy. Depending upon the target cells and therapeutic nucleic acid, the present invention can be used in treatment of various conditions and diseases, to
 5 increase expression of a desired protein, to inhibit expression or function of a gene product, and so forth. For instance, transcription or translation of a given therapeutic nucleic acid molecule may be useful in treating cancer or an acquired disease, such as AIDS, pneumonia, emphysema, or in correcting inborn errors of metabolism, such as cystic fibrosis. Transcription or translation of a therapeutic nucleic acid may also
 10 effect contraceptive sterilization, including contraceptive sterilization of feral animals. Allergen-mediated and infectious agent-mediated inflammatory disorders also can be countered by administering, via the present invention, a therapeutic nucleic acid molecule that, upon expression in a patient, affects immune response(s) associated with the allergen and infectious agent, respectively. A therapeutic nucleic acid
 15 molecule also may have an expression product, or there may be a downstream product of post-translational modification of the expression product, that reduces the immunologic sequelae related to transplantation or that helps facilitate tissue growth and regeneration.

The invention also relates to the transfer of nucleic acids into selected cell
 20 types *in vitro*. Such transfers are useful for a variety of purposes, such as to create a cell that can produce large quantities of a selected protein, which can then be harvested.

In a related aspect, the invention provides a composition of matter useful for introducing exogenous nucleic acid molecules into target non-phagocytic mammalian
 25 cells with high efficiency. The composition comprises (i) a bacterially derived minicell and (ii) a bispecific ligand. The minicell and ligand may be any of those described herein. Thus, the minicell contains a therapeutic nucleic acid molecule and the bispecific ligand preferably is capable of binding to a surface component of the minicell and to a surface component of a target mammalian cell.

A composition consisting essentially of recombinant minicells and bispecific ligands of the present invention (that is, a composition that includes such minicells and ligands with other constituents that do not interfere unduly with the DNA-delivering quality of the composition) can be formulated in conventional manner, using one or
5 more physiologically acceptable carriers or excipients. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or vials, or in multi-dose containers, with or without an added preservative. The formulation can be a solution, a suspension, or an emulsion in oily or aqueous vehicles, and may contain formulatory agents, such as suspending, stabilizing and/or dispersing agents. A suitable solution is isotonic with the
10 blood of the recipient and is illustrated by saline, Ringer's solution, and dextrose solution. Alternatively, compositions may be in lyophilized powder form, for reconstitution with a suitable vehicle, *e.g.*, sterile, pyrogen-free water or physiological saline. The compositions also may be formulated as a depot preparation. Such long-acting formulations may be administered by implantation (for example, subcutaneously
15 or intramuscularly) or by intramuscular injection.

A composition of the present invention can be administered via various routes and to various sites in a mammalian body, to achieve the therapeutic effect(s) desired, either locally or systemically. Delivery may be accomplished, for example, by oral administration, by application of the formulation to a body cavity, by inhalation or
20 insufflation, or by parenteral, intramuscular, intravenous, intraportal, intrahepatic, peritoneal, subcutaneous, intratumoral, or intradermal administration. The mode and site of administration is dependent on the location of the target cells. For example, cystic-fibrotic cells may be efficiently targeted by inhaled delivery of the targeted recombinant minicells. Similarly, tumor metastasis may be more efficiently treated
25 via intravenous delivery of targeted recombinant minicells. Primary ovarian cancer may be treated via intraperitoneal delivery of targeted recombinant minicells.

The following examples are intended to illustrate and provide a more complete understanding of the invention without limiting the invention to the examples provided.

Example 1. Highly efficient binding and receptor-mediated internalization of bispecific antibody-targeted minicells into non-phagocytic human prostate carcinoma cells.

5 This experiment demonstrates that a bispecific antibody with Fab fragments carrying anti-*S. typhimurium* LPS and anti-androgen receptor binding specificities can enable binding and receptor-mediated internalization of *S. typhimurium*-derived minicells into prostate carcinoma cells that are known to over-express the androgen receptor on the cell surface.

S. typhimurium minCDE- mutant strain generated previously (patent application, PCT/IB02/04632) was transformed with recombinant plasmid pORF5-
10 HSV1tk::Sh ble (Invivogen, San Diego, CA, USA). The plasmid is a mammalian gene expression vector that expresses the HSV1tk::Sh ble fusion gene under the control of the EF-1 α / eIF4g hybrid promoter. The HSV1tk is a suicide gene from Herpes simplex serotype 1 virus (HSV1) and encodes an enzyme, thymidine kinase,
15 that can convert prodrug guanosine analog ganciclovir (GCV) to ganciclovir-monophosphate (GCV-MP). The latter is then converted to the diphosphate and triphosphate forms by endogenous kinases. GCV-triphosphate lacks the 3' OH on the deoxyribose as well as the bond between the 2' and 3' carbons which are necessary for DNA chain elongation. As a result, GCV-triphosphate integration causes
20 premature DNA chain termination and leads to apoptosis. Expression of HSV1tk therefore sensitizes transfected mammalian cells to ganciclovir and is one of the most widely used single suicide strategies for cancer gene therapy (Singhal and Kaiser, 1998). As a control, a plasmid was constructed where HSVtk::Sh ble gene fusion was deleted by cleaving plasmid pORF5-HSV1tk::Sh ble with restriction enzymes *NcoI*
25 and *NheI*, blunt-ending the sites with T4 DNA polymerase and religating the plasmid. The *NcoI* and *NheI* sites are unique in plasmid pORF5-HSV1tk::Sh ble and flank the HSV1tk::Sh ble gene fusion. The resulting plasmid designated pORF5-HSV1tk- was also transformed in *S. typhimurium minCDE*- mutant strain.

Recombinant minicells carrying the plasmids were purified using the gradient centrifugation / filamentation / filtration / endotoxin removal procedure described in patent application, PCT/IB02/04632 and U.S. patent application number 10/602.021.

The bispecific antibody was constructed by linking anti-*S. typhimurium* lipopolysaccharide (Biodesign, Saco, Maine, USA) and anti-androgen receptor mouse monoclonal antibodies (IgG; Abcam, Cambridge, UK) to purified recombinant protein A/G via the Fc fragments of each monoclonal antibody and in brief the procedure was as follows.

Purified recombinant protein A/G (Pierce Biotechnology, Rockford, IL, USA) was diluted to a final concentration of 100 ug/ml in Immunopure binding buffer (Pierce Biotechnology) and 0.5 ml of the solution was incubated overnight at 4°C with a premixed solution containing 20 ug/ml each of anti-*S. typhimurium* LPS (Research Diagnostics Inc., Flanders, NJ, USA) and anti-human androgen receptor (Abcam, Cambridge, UK) monoclonal antibodies. The excess antibodies unbound to protein A/G were then removed as follows. Dynabeads® Protein G solution (Dynabeads® [2.8 µm] coated with recombinant Protein G covalently coupled to the surface of the magnetic particles; Dynal Biotech, Oslo, Norway) was mixed gently and 100 µl of the solution was transferred into an eppendorf centrifuge tube. The tube was placed in the Dynal MPC-S (Magnetic Particle Concentrator, type S) to immobilize the beads and the supernatant was discarded. The beads were resuspended in 0.5 ml of washing solution containing 0.1M Na-phosphate buffer (pH 5.0). The bead immobilization and washing steps were repeated three times. The solution containing protein A/G-bispecific antibody complex was added to the beads and incubated with gentle mixing at room temperature for 40 min. The tube was placed on the MPC-S stand to immobilize the beads and the protein A/G-bispecific antibody complex was removed with a pipette. This step removed the unbound excess monoclonal antibodies from the solution and provided a solution that carried the bispecific antibody linked to protein A/G via their Fc fragments.

10¹⁰ recombinant minicells were incubated with the protein A/G-bispecific antibody for 1 hr at room temperature to coat the minicells with the antibody via its anti-LPS Fab region.

Prostate carcinoma cells, LNCaP (ATCC, Rockville, MD, USA) were grown to full confluency in T-75 flasks in RPMI 1640 medium supplemented with 10% FCS and antibiotics. The cells were passaged in T-25 flasks at 50% confluency. After overnight attachment, the culture medium was refreshed and to one flask was added 10⁷ recombinant minicells carrying plasmid pORF5-HSV1tk::Sh ble (non-targeted recombinant minicells) and to another flask was added 10⁷ of the same minicells but carrying cell surface attached bispecific antibody (targeted recombinant minicells). The ratio of minicells to prostate carcinoma cells was 100:1. The transfected cells were incubated in an incubator under 5% CO₂ and 37°C for 16, 24 and 36 hrs followed by four washes (5ml per wash) with fresh 1x Dulbecco's medium with gentle shaking. All cells were trypsinized and then passaged on 13mm coverslips in 24 well plate (each time point in triplicate), with cell numbers in sub-confluency.

The cells on coverslips were fixed with 4% paraformaldehyde for 30mins and blocked with 5% normal goat serum overnight followed by staining with anti-*S. typhimurium* LPS (1:200; Biodesign, Saco, Maine, USA) monoclonal antibody. The antibody binding was revealed with goat anti-mouse IgG conjugated with Alexa Fluor 594 (1:1000, red fluorescence; excitation 590nm and emission 617nm; Molecular Probes, Eugene, OR, USA) and viewed by fluorescence confocal microscopy (Fluoview, Olympus America, Melville, NY, USA). Fluorescence and Differential Image Contrast (DIC) images were collected and overlaid as shown in Figure 1.

The results showed that non-targeted recombinant minicells did not specifically adhere to or get internalized in the LNCaP prostate carcinoma cells at any of the time points analyzed (Fig. 1B and 1D) and cells appeared the same as control non-transfected cells. All fields analyzed revealed minor background red fluorescence. In contrast, the targeted recombinant minicells were found to strongly adhere to the LNCaP cells presumably via binding of the targeting bispecific antibody

to the cell surface androgen receptor. Additionally, at the 16hr and 24hr incubation time points, most LNCaP cells showed intense red fluorescence within the cytoplasm of the cells (Fig. 1C, 1E and 1F) indicating that the minicells had been internalized via receptor-mediated endocytosis.

5 This result suggested that the minicells carrying surface-attached bispecific antibody mediated highly efficient binding of the minicells to the cell surface receptor found on a mammalian cell (androgen receptor in the above example) and that the adherent minicells were rapidly internalized by the non-phagocytic mammalian cell (prostate carcinoma cell in the above example).

10 **Example 2. Highly efficient binding and receptor-mediated internalization of bispecific antibody-targeted minicells into non-phagocytic human breast adenocarcinoma cells.**

Example 1 demonstrated that a bispecific antibody with anti-LPS (minicell specificity) and anti-androgen receptor binding specificity can efficiently enable
15 strong binding to the androgen receptor on a non-phagocytic mammalian cell, the prostate carcinoma cell. Additionally, the results demonstrated that the receptor binding triggered receptor-mediated endocytosis of the recombinant minicells at a high efficiency. This example demonstrates that the above-observed phenomenon is generalized and that the invention and discovery was applicable to a range of different
20 endocytosis-competent receptors on different non-phagocytic mammalian cells.

Accordingly, this experiment demonstrates that human breast adenocarcinoma cells (MDA-MB-468, ATCC; human mammary epithelial cells; non-phagocytic) can be targeted via a bispecific antibody carrying Fab fragments with anti-*S. typhimurium* LPS (minicell surface binding specificity) and anti-epidermal growth factor receptor
25 (EGFR) binding specificity. The cell line MDA-MB-468 cells were grown in tissue culture as described for prostate carcinoma cells in example 1. The bispecific antibody was constructed as described in Example 1, except that the anti-androgen receptor monoclonal antibody was replaced with anti-EGFR monoclonal antibody (Oncogene Research Products, Cambridge, MA, USA). Targeted and non-targeted

recombinant minicells were generated and used to transfect the MDA-MB-468 cells and the cells were stained for *S. typhimurium* LPS (minicells) at time intervals of 16hrs, 24hrs and 36hrs as described above for prostate carcinoma cells.

The results revealed (Fig. 2) that control cells and cells treated with non-targeted minicells exhibited only minor background red fluorescence at all the time points (Figs 2A and 2B), suggesting that the minicells were unable to adhere to and transfect the non-phagocytic mammalian cells. In contrast, the cells treated with targeted minicells exhibited strong red fluorescence in the cytoplasm after 24hrs incubation and the fluorescence increased to cover more of the cytoplasm after 36hrs (Figs 2C-E). This suggested that the bispecific antibody enabled the strong binding of the minicells to the EGF receptor on the surface of MDA-MB-468 cells and that the binding triggered receptor mediated endocytosis of the minicells.

Example 3. Highly efficient binding and receptor-mediated internalization of bispecific antibody-targeted minicells into non-phagocytic human ovarian carcinoma cells.

Examples 1 and 2 demonstrated that a bispecific antibody with anti-LPS (minicell specificity) and either anti-androgen receptor binding specificity or anti-EGFR specificity can efficiently enable strong binding to the androgen receptor or EGFR on a non-phagocytic prostate carcinoma cells and breast carcinoma cells respectively. Additionally, the results demonstrated that the receptor binding triggered receptor-mediated endocytosis of the recombinant minicells at a high efficiency. This example further demonstrates the general applicability of the invention and discovery.

Accordingly, this experiment demonstrates that human ovarian carcinoma cells (SKOV-3, ATCC; epithelial cells; non-phagocytic) can be targeted via a bispecific antibody carrying Fab fragments with anti-*S. typhimurium* LPS (minicell surface binding specificity) and mouse anti-human Her2/neu receptor (Serotec Inc., Raleigh, NC, USA) binding specificity. SKOV-3 cells are known to overexpress the Her2 receptor (Salomon *et al.*, 1995). The experiment was performed as described in

Examples 1 and 2, and the samples were stained for anti-LPS (red fluorescence) as before.

The results (Fig. 3) were similar to those obtained in examples 1 and 2. The control SKOV-3 cells and those treated with non-targeted minicells, showed only
5 minor background red fluorescence.

Example 4. Bispecific antibody-mediated targeting of minicells to human breast cancer xenografts in female athymic nude mice.

This example demonstrates that targeted recombinant minicells carrying a plasmid encoding HSVtk gene can effect regression of human breast cancer cell
10 tumor xenografts established in 6 week old female athymic nude mice.

The bispecific antibody was constructed as described in Example 1, except that instead of the anti-androgen receptor monoclonal antibody, the anti-epidermal growth factor receptor (anti-EGFR) monoclonal antibody (Oncogene Research Products, Cambridge, MA, USA) was used. This was because the xenografted cells
15 were human breast cancer cells MDA-MB-468 that are known to overexpress the EGF receptor on the cell surface. The mice were purchased from Animal Resources Centre, Perth, WA, and all animal experiments were performed in compliance with the guide of care and use of laboratory animals and with Animal Ethics Committee approval. The experiments were performed in the NSW Agriculture accredited small
20 animal facility at EnGeneIC Pty Ltd (Sydney, NSW, Australia). MDA-MB-468 human breast cancer cells were cultured as described in example 2 and 1.5×10^6 cells in 50 uL serum-free media together with 50 uL growth factor reduced matrigel (BD Biosciences, Franklin Lakes, NJ, USA) were injected subcutaneously between the shoulder blades of each mouse using a 23-gauge needle. The tumors were measured
25 twice a week using an electronic digital caliper (Mitutoyo, Japan, precision to 0.001) and tumor volume was calculated using the formula, length (mm) x width² (mm) X 0.5 = volume (mm³). 21 days post-implantation the tumors reached volumes between 50 mm³ and 80 mm³, and mice were randomized to six different groups of 12 per group.

The experiment was designed as follows. Group 1 (control) received no treatment. Group 2 (control) received non-targeted recombinant minicells that carried plasmid pORF5- HSV1tk::Sh ble (designated M-HSVtk) on days 21, 28 and 35. The mice also received GCV on days 25, 26, 32, 33, 39 and 40 i.e. 2 doses of GCV on successive days. This group was designed to determine if non-targeted minicells could deliver the suicide gene to the tumor cells and affect tumor regression following GCV treatment. Group 3 (control) was designed to determine if treatment with targeted recombinant minicells carrying plasmid pORF5-HSV1tk::Sh ble in the absence of GCV had any effect on tumor regression. Therefore, Group 3 mice received targeted recombinant minicells carrying plasmid pORF5-HSV1tk::Sh ble (designated TM-HSVtk) on the same days as for group 2 but received no GCV treatment. Group 4 (control) was designed to determine if the bispecific antibody in the absence of recombinant minicells had any effect on tumor regression. Therefore these mice received the bispecific antibody on the same days that recombinant targeted or non-targeted minicells were given, i.e., days 21, 28 and 35. The antibody treatment was followed by GCV treatment on the same days as for group 2. Group 5 (experimental) was designed to determine if the targeted recombinant minicells carrying plasmid pORF5-HSV1tk::Sh ble could effectively deliver the plasmid to the xenografted tumor cells and if tumor regression could be observed following treatment of the mice with a single dose of GCV after each minicell dose. Therefore, group 5 received targeted recombinant minicells on the same days as for group 3 followed by GCV treatment on days 25, 33 and 39. Group 6 (experimental) was the same as group 5 but received 2 doses of GCV on successive days as for groups 2 and 4.

Mice receiving the respective minicells were injected intratumorally with 10^8 minicells resuspended in 30 ul of sterile physiological saline. Gene targeting experiments *in-vitro* in MDA-MB-468 cells had revealed that the minicell delivered plasmid expressed the HSVtk enzyme after at least 48hrs post-transfection with the targeted recombinant minicells. Therefore, groups 2, 4, 5 and 6 were given GCV after 3 to 4 days post-minicell inoculation to allow the transfected tumor xenograft cells to

sufficiently express the HSVtk enzyme to be responsive to GCV. GCV was administered intraperitoneally at a concentration of 100 mg/kg of mouse weight.

Figure 4 shows the progression in tumor volume over the course of the experiment. The results revealed that only targeted recombinant minicells (Groups 5 and 6) were able to successfully deliver the HSV1tk gene encoding plasmid to the xenografted tumor cells. The tumor volumes in these two groups did not increase in size and remained stable throughout the course of the experiment. In contrast, the tumor volumes rapidly increased in the four control groups (Groups 1-4). Interestingly, group 2 mice also showed no evidence of tumor regression, suggesting the non-targeted recombinant minicells could not transfect the human breast cancer cells. Statistical analysis of the data using One-way ANOVA showed that experimental groups (5 and 6) were highly significant compared to the control groups 1 to 4 ($p=0.001$). This result is a first demonstration of targeted *in-vivo* gene delivery to non-phagocytic mammalian cells mediated by bacterially derived intact recombinant minicells. It also demonstrates a role for receptor-mediated endocytosis of the minicells in achieving highly significant gene delivery to these non-phagocytic mammalian cells (compare group 2 with groups 5 and 6).

CITED PUBLICATIONS

This application incorporates by reference each of the following publications:

Batra RK, Wang-Johanning F, Wagner E, Garver RI Jr, Curiel DT. Receptor-mediated gene delivery employing lectin-binding specificity. *Gene Ther.* 1994 Jul;1(4):255-60.

Becker CM, Farnebo FA, Iordanescu I, Behonick DJ, Shih MC, Dunning P, Christofferson R, Mulligan RC, Taylor GA, Kuo CJ, Zetter BR. Gene therapy of prostate cancer with the soluble vascular endothelial growth factor receptor Flk1. *Cancer Biol Ther.* 2002 Sep-Oct;1(5):548-53.

Bondoc, LL and Fitzpatrick S. Size distribution analysis of recombinant adenovirus using disc centrifugation. *J Indust Micro Biotechnol.* 20: 317-322 (1998).

Britton *et al.*, "Characterization of a prokaryotic SMC protein involved in chromosome partitioning," *Genes Dev.* 12: 1254 (1998).

Carter, P. Improving the efficacy of antibody-based cancer therapies. *Nat Rev Cancer.* 2001 Nov;1(2):118-29.

- 5 Ciliberto *et al.*, "Cell-specific expression of a transfected human alpha 1-antitrypsin gene," *Cell.* 41: 531 (1985).

Conner SD, Schmid SL. Regulated portals of entry into the cell. *Nature.* 2003 Mar 6;422(6927):37-44.

- 10 Curiel *et al.*, "Long-term inhibition of clinical and laboratory human immunodeficiency virus strains in human T-cell lines containing an HIV-regulated diphtheria toxin A chain gene," *Hum. Gene Ther.* 4: 741 (1993).

de Boer *et al.*, "Roles of MinC and MinD in the site-specific septation block mediated by the MinCDE system of *Escherichia coli*," *J. Bacteriol.* 174: 63 (1992).

- 15 de Haard, H. J. *et al.* A large non-immunized human Fab fragment phage library that permits rapid isolation and kinetic analysis of high affinity antibodies. *J. Biol. Chem.* 274, 18218-18230 (1999).

Dinges *et al.*, "HIV-regulated diphtheria toxin A chain gene confers long-term protection against HIV type 1 infection in the human promonocytic cell line U937," *Hum. Gene Ther.* 6: 1437 (1995).

- 20 Dubel S, Breitling F, Kontermann R, Schmidt T, Skerra A, Little M. Bifunctional and multimeric complexes of streptavidin fused to single chain antibodies (scFv). *J. Immunol. Methods* (1995) 178, 201-209.

Fernandez T, Bayley H. 1998. Ferrying proteins to the other side. *Nat Biotechnol* 16:418-430.

Frain *et al.*, "Binding of a liver-specific factor to the human albumin gene promoter and enhancer," *Mol. Cell Biol.* 10: 991 (1990).

Glennie MJ, McBride HM, Worth AT, Stevenson GT. Preparation and performance of bispecific F(ab' gamma)2 antibody containing thioether-linked Fab' gamma fragments. *J Immunol.* 1987 Oct 1;139(7):2367-75.

Gosselin MA, Lee RJ. Folate receptor-targeted liposomes as vectors for therapeutic agents. *Biotechnol Annu Rev.* 2002;8:103-31.

Griffiths, A. D. *et al.* Isolation of high affinity human antibodies directly from large synthetic repertoires. *EMBO J.* 13, 3245-3260 (1994).

10 Grillot-Courvalin C, Goussard S, Courvalin P. Wild-type intracellular bacteria deliver DNA into mammalian cells. *Cell Microbiol.* 2002 Mar; 4(3):177-86.

Harrison *et al.*, "Inhibition of human immunodeficiency virus-1 production resulting from transduction with a retrovirus containing an HIV-regulated diphtheria toxin A chain gene," *Hum. Gene Ther.* 3: 461 (1992a).

15 Harrison *et al.*, "Inhibition of HIV production in cells containing an integrated, HIV-regulated diphtheria toxin A chain gene," *AIDS Res. Hum. Retroviruses* 8: 39 (1992b).

Harry, "Bacterial cell division: Regulating Z-ring formation," *Mol. Microbiol.* 40: 795 (2001).

20 Hart, "Tissue specific promoters in targeting systematically delivered gene therapy," *Semin. Oncol.* 23: 154 (1996).

Heim *et al.*, "Wavelength mutations and posttranslational autooxidation of green fluorescent protein," *Proc. Nat'l. Acad. Sci. USA* 91: 12501 (1994).

Hiraga *et al.*, "Chromosome partitioning in *Escherichia coli*: novel mutants producing anucleate cells," *J. Bacteriol.* 171: 1496 (1989).

Hoshida T, Sunamura M, Duda DG, Egawa S, Miyazaki S, Shineha R, Hamada H, Ohtani H, Satomi S, Matsuno S. Gene therapy for pancreatic cancer using an adenovirus vector encoding soluble flt-1 vascular endothelial growth factor receptor. *Pancreas*. 2002 Aug;25(2):111-21.

- 5 Hu, S, L Shively, A Raubitschek, M Sherman, LE Williams, JY Wong, JE Shively, and AM Wu. Minibody: A novel engineered anti-carcinoembryonic antigen antibody fragment (single-chain Fv-CH3) which exhibits rapid, high-level targeting of xenografts *Cancer Res*. 1996 56: 3055-3061.

- 10 Hu & Lutkenhaus, "Topological regulation of cell division in *Escherichia coli* involves rapid pole to pole oscillation of the division inhibitor MinC under the control of MinD and MinE," *Mol. Microbiol*. 34: 82 (1999).

Hudson, P.J. & Souriau, C. Recombinant antibodies for cancer diagnosis and therapy. *Expert Opin. Biol. Ther.* 1, 845-855 (2001).

Hudson PJ, Souriau C. Engineered antibodies. *Nat Med*. 2003 Jan;9 (1):129-34.

- 15 - Hung MC, Hortobagyi GN, Ueno NT. Development of clinical trial of E1A gene therapy targeting HER-2/neu-overexpressing breast and ovarian cancer. *Adv Exp Med Biol*. 2000;465:171-80.

Ireton *et al.*, "spo0J is required for normal chromosome segregation as well as the initiation of sporulation in *Bacillus subtilis*," *J. Bacteriol*. 176: 5320 (1994).

- 20 Jones, P. T., Dear, P. H., Foote, J., Neuberger, M. S. & Winter, G. Replacing the complementarity-determining regions in a human antibody with those from a mouse. *Nature* 321, 522-525 (1986).

- Katabi *et al.*, "Hexokinase Type II: A Novel Tumor Specific Promoter for Gene-Targeted Therapy Differentially Expressed and Regulated in Human Cancer Cells,"
25 *Human Gene Therapy* 10: 155 (1999).

Kaetzel CS, Blanch VJ, Hempen PM, Phillips KM, Piskurich JF, Youngman KR
(1997): The polymeric immunoglobulin receptor: structure and synthesis. *Biochem
Soc Trans* 25:475-480.

Karpovsky B, Titus JA, Stephany DA, Segal DM. Production of target-specific
5 effector cells using hetero-cross-linked aggregates containing anti-target cell and anti-
Fc gamma receptor antibodies. *J Exp Med.* 1984 Dec 1;160(6):1686-701.

Kelsey *et al.*, "Species- and tissue-specific expression of human alpha 1-antitrypsin in
transgenic mice," *Genes and Devel.* 1: 161 (1987).

Kerem *et al.*, "Identification of the cystic fibrosis gene: genetic analysis," *Science*
10 245: 1073 (1989).

Khare, P. D. *et al.* Tumor growth suppression by a retroviral vector displaying scFv
antibody to CEA and carrying the iNOS gene. *Anticancer Res.* 22, 2443-2446 (2002).

Kleeff J, Fukahi K, Lopez ME, Friess H, Buchler MW, Sosnowski BA, Korc M.
Targeting of suicide gene delivery in pancreatic cancer cells via FGF receptors.
15 *Cancer Gene Ther.* 2002 Jun;9(6):522-32.

Knappik, A. *et al.* Fully synthetic human combinatorial antibody libraries (HuCAL)
based on modular consensus frameworks and CDRs randomized with trinucleotides.
J. Mol. Biol. 296, 57-86 (2000).

Kostelny SA, Cole MS, Tso JY. Formation of a bispecific antibody by the use of
20 leucine zippers. *J Immunol.* 1992 Mar 1;148(5):1547-53.

Kurane *et al.*, "Targeted Gene Transfer for Adenocarcinoma Using a Combination of
Tumor specific Antibody and Tissue-specific Promoter," *Jpn. J. Cancer Res.* 89: 1212
(1998).

Levin *et al.*, "Identification of *Bacillus subtilis* genes for septum placement and shape
25 determination," *J. Bacteriol.* 174: 6717 (1992).

Leder *et al.*, "Consequences of widespread deregulation of the c-myc gene in transgenic mice: multiple neoplasms and normal development," *Cell* 45: 485 (1986).

MacDonald *et al.*, "Expression of the pancreatic elastase I gene in transgenic mice," *Hepatology* 7: 425 (1987).

- 5 Mason *et al.*, "The hypogonadal mouse: reproductive functions restored by gene therapy," *Science* 234: 1372 (1986).

Morton & Potter, "Rhabdomyosarcoma-specific expression of the herpes simplex virus thymidine kinase gene confers sensitivity to ganciclovir," *J. Pharmacology & Exper. Therapeutics* 286: 1066 (1998).

- 10 Okada *et al.*, "Possible function of the cytoplasmic axial filaments in chromosomal segregation and cellular division of *Escherichia coli*," *Sci. Prog.* 77: 253 (1993-94).

Okada *et al.*, "Cytoplasmic axial filaments in *Escherichia coli* cells: possible function in the mechanism of chromosome segregation and cell division," *J. Bacteriol.* 176: 917 (1994).

- 15 Osbourn, J., Jermutus, L., Duncan, A. Current methods for the generation of human antibodies for the treatment of autoimmune diseases. *Drug Delivery Tech* 8: 845-851 (2003).

Pack P, Pluckthun A. Miniantibodies: use of amphipathic helices to produce functional, flexibly linked dimeric Fv fragments with high avidity in *Escherichia coli*.

- 20 *Biochemistry*. 1992 Feb 18;31(6):1579-84.

Phelan A, Elliott G, O'Hare P. 1998. Intercellular delivery of functional p53 by the herpesvirus protein VP22. *Nat Biotechnol* 16:440-443.

Pinkert *et al.*, "An albumin enhancer located 10 kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice," *Genes and*

- 25 *Devel.* 1: 268 (1987).

- Prasher *et al.*, "Using GFP to see the light," *Trends in Genetics* 11: 320 (1995).
- Ragheb *et al.*, "Inhibition of human immunodeficiency virus type 1 by Tat/Rev-regulated expression of cytosine deaminase, interferon alpha2, or diphtheria toxin compared with inhibition by transdominant Rev," *Hum. Gene Ther.* 10: 103 (1999).
- 5 Raskin & de Boer, "MinDE-dependent pole-to-pole oscillation of division inhibitor MinC in *Escherichia coli*," *J. Bacteriol.* 181: 6419 (1999).
- Readhead *et al.*, "Myelin deficient mice: expression of myelin basic protein and generation of mice with varying levels of myelin," *Cell* 48: 703 (1987).
- Reeve, "Use of minicells for bacteriophage-directed polypeptide synthesis," *Methods*
10 *Enzymol.* 68: 493 (1979).
- Ridgway JB, Presta LG, Carter P. 'Knobs-into-holes' engineering of antibody CH3 domains for heavy chain heterodimerization. *Protein Eng.* 1996 Jul;9(7):617-21.
- Riechmann, L., Clark, M., Waldmann, H. & Winter, G. Reshaping human antibodies for therapy. *Nature* 332, 323-327 (1988).
- 15 Riordan *et al.*, "Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA," *Science* 245: 1066 (1989).
- Rojas M, Donahue JP, Tan Z, Lin YZ. 1998. Genetic engineering of proteins with cell membrane permeability. *Nat Biotechnol* 16(4): 370-375.
- Rommens *et al.*, "Identification of the cystic fibrosis gene: Chromosome walking and
20 jumping," *Science* 245: 1059 (1989).
- Salomon DS, Brandt R, Ciardiello F, Normanno N. Epidermal growth factor-related peptides and their receptors in human malignancies. *Crit Rev Oncol Hematol* 1995, 19, 183-232.
- Shangara *et al.*, "Suicide genes: past, present and future perspectives," *Immunology*
25 *Today* 21: 48 (2000).

- Sheets, M. D. *et al.* Efficient construction of a large nonimmune phage antibody library: the production of high-affinity human single-chain antibodies to protein antigens. *Proc. Natl Acad. Sci. USA* 95, 6157–6162 (1998).
- 5 Spencer, “Developments in suicide genes for preclinical and clinical applications,” *Molecular Therapeutics* 2: 433 (2000).
- Stewart & D'Ari, “Genetic and morphological characterization of an *Escherichia coli* chromosome segregation mutant,” *J. Bacteriol.* 174: 4513 (1992).
- Swift *et al.*, “Tissue-specific expression of the rat pancreatic elastase I gene in transgenic mice,” *Cell* 38: 639 (1984).
- 10 Thomas, C. E., Edwards, P., Wickham, T. J., Castro, M. G. & Lowenstein, P. R. Adenovirus binding to the coxsackievirus and adenovirus receptor or integrins is not required to elicit brain inflammation but is necessary to transduce specific neural cell types. *J. Virol.* 76, 3452–3460 (2002).
- 15 Thomas CE, Ehrhardt A, Kay MA. Progress and problems with the use of viral vectors for gene therapy. *Nat Rev Genet.* 2003 May;4(5):346-58.
- Thurnher M, Wagner E, Clausen H, Mechtler K, Rusconi S, Dinter A, Birnstiel ML, Berger EG, Cotten M. Carbohydrate receptor-mediated gene transfer to human T leukaemic cells. *Glycobiology.* 1994 Aug;4(4):429-35.
- 20 Todorovska, A. *et al.* Design and application of diabodies, triabodies and tetrabodies for cancer targeting. *J. Immunol. Methods* 248, 47–66 (2001).
- Tomlinson, I. & Holliger, P. Methods for generating multivalent and bispecific antibody fragments. *Methods Enzymol.* 326, 461–479 (2000).
- Vaughan, T. J. *et al.* Human antibodies with subnanomolar affinities isolated from a large non-immunized phage display library. *Nature Biotechnol.* 14, 309–314 (1996).

Vaughan, T. J., Osbourn, J. K. & Tempest, P. R. Human antibodies by design. *Nature Biotechnol.* 16, 535–539 (1998).

Verhoeyen, M., Milstein, C. & Winter, G. Reshaping human antibodies: grafting an antilysozyme activity. *Science* 239, 1534–1536 (1988).

- 5 Wachi *et al.*, “New mre genes *mreC* and *mreD*, responsible for formation of the rod shape of *Escherichia coli* cells,” *J. Bacteriol.* 171: 6511 (1989).

Yazawa *et al.*, “Current progress in suicide gene therapy for cancer,” *World J. Surg.* 26: 783 (2002).

Ziady AG, Perales JC, Ferkol T, Gerken T, Beegen H, Perlmutter DH, Davis PB.

- 10 Gene transfer into hepatoma cell lines via the serpin enzyme complex receptor. *Am J Physiol.* 1997 Aug;273(2 Pt 1):G545-52.

WHAT IS CLAIMED IS:

1. A targeted gene delivery method that comprises bringing bispecific ligands into contact with (a) bacterially derived minicells that contain a therapeutic nucleic acid sequence and (b) non-phagocytic mammalian cells, such that (i) said
5 bispecific ligands cause said minicells to bind to said mammalian cells and (ii) said minicells are engulfed by said mammalian cells, which produce an expression product of said therapeutic nucleic acid sequence.
2. A method according to claim 1, wherein said bispecific ligand comprises polypeptide or carbohydrate.
- 10 3. A method according to claim 1, wherein said bispecific ligand comprises a first arm that carries specificity for a bacterially derived minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor.
4. A method according to claim 3, wherein said first arm and said second
15 arm are monospecific.
5. A method according to claim 3, wherein said first arm and said second arm are multivalent.
6. A method according to claim 3, wherein said minicell surface structure is an O-polysaccharide component of a lipopolysaccharide on said minicell surface.
- 20 7. A method according to claim 3, wherein said minicell surface structure is a member of the group consisting of outer membrane proteins, pilli, fimbriae, flagella, and cell-surface exposed carbohydrates.
8. A method according to claim 3, wherein said mammalian cell surface receptor is capable of activating receptor-mediated endocytosis of said minicell.

9. A method according to claim 1, wherein said bispecific ligand comprises an antibody or antibody fragment.

10. A method according to claim 1, wherein said bispecific ligand comprises a humanized antibody.

5 11. A method according to claim 1, wherein said minicell comprises an intact cell wall.

12. A method according to claim 1, wherein said therapeutic nucleic acid sequence encodes a suicide gene.

10 13. A method according to claim 1, wherein said therapeutic nucleic acid encodes a normal counterpart of a gene that expresses a protein that functions abnormally or is present in abnormal levels in said mammalian cells.

14. A method according to claim 1, wherein said mammalian cells are *in vitro*.

15 15. A method according to claim 1, wherein said mammalian cells are *in vivo*.

16. A method according to claim 1, wherein said therapeutic nucleic acid is contained on a plasmid comprised of multiple nucleic acid sequences.

17. A method according to claim 16, wherein said plasmid comprises a regulatory element.

20 18. A method according to claim 16, wherein said plasmid comprises a reporter element

25 19. A composition comprising (i) a bacterially derived minicell that contains a therapeutic nucleic acid molecule and (ii) a bispecific ligand that is capable of binding to a surface component of said minicell and to a surface component of a non-phagocytic mammalian cell.

20. Use of bacterially derived intact minicells and bispecific ligands in the preparation of a medicament, said minicells containing a therapeutic nucleic acid molecule and said bispecific ligands being capable of binding to said minicells and to target non-phagocytic mammalian cells, for use in a method of treating a disease or
5 modifying a trait by administration of said medicament to a cell, tissue, or organ.

ABSTRACT

A method of targeting bacterially derived intact minicells to specific non-phagocytic mammalian cells employs bi-specific ligands to efficiently deliver nucleic acids to the mammalian cells. Bispecific ligands comprising a first arm that carries
5 specificity for a bacterially derived minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor are useful for targeting minicells to specific non-phagocytic mammalian cells and causing endocytosis of minicells by non-phagocytic cells.

10

Title: METHODS FOR TARGETED GENE
DELIVERY TO NON-PHAGOCYTIC
MAMMALIAN CELLS VIA BACTERIALLY
DERIVED INTACT MINICELLS IN-VITRO AND
IN-VIVO

Inventor(s): Himanshu BRAHMBHATT et al.
DOCKET NO.: 060348-0105

Figure 1





Figure 2


A
Controls, MDA-MB-468 alone




B
MDA-MB-468, non-targeting (24hrs)




C
MDA-MB-468, targeting (24hrs)



D
MDA-MB-468, targeting (24hrs)



E
MDA-MB-468, targeting (36hrs)



Title: METHODS FOR TARGETED GENE
DELIVERY TO NON-PHAGOCYTIC
MAMMALIAN CELLS VIA BACTERIALLY
DERIVED INTACT MINICELLS IN-VITRO AND
IN-VIVO
Inventor(s): Himanshu BRAHMBHATT et al.
DOCKET NO.: 060348-0105

Figure 3





Figure 4



Application Data Sheet

Application Information

Application Type::	Provisional
Subject Matter::	Utility
Suggested classification::	
Suggested Group Art Unit::	
CD-ROM or CD-R?::	None
Computer Readable Form (CRF)?::	No
Title::	METHODS FOR TARGETED GENE DELIVERY TO NON-PHAGOCYTIC MAMMALIAN CELLS VIA BACTERIALLY DERIVED INTACT MINICELLS <i>IN- VITRO</i> AND <i>IN-VIVO</i> 060348-0105
Attorney Docket Number::	
Request for Early Publication?::	No
Request for Non-Publication?::	No
Suggested Drawing Figure::	1
Total Drawing Sheets::	4
Small Entity?::	Yes
Petition included?::	No
Secrecy Order in Parent Appl.?::	No

Applicant Information

Applicant Authority Type::	Inventor
Primary Citizenship Country::	US
Status::	Full Capacity
Given Name::	Himanshu
Family Name::	BRAHMBHATT
City of Residence::	Sydney
Country of Residence::	
Street of mailing address::	96 Prairie Vale Road Bossley Park Sydney NSW 2176
Country of mailing address::	Australia
Applicant Authority Type::	Inventor
Primary Citizenship Country::	Australian
Status::	Full Capacity

Given Name:: Jennifer
Family Name:: MACDIARMID
City of Residence:: Sydney
Country of Residence:: Australia
Street of mailing address:: 8 Lombard Street Glebe
Sydney
NSW 2037
Country of mailing address:: Australia

Correspondence Information

Correspondence Customer Number:: 22428
E-Mail address:: PTOMailWashington@Foley.com

Representative Information

Representative Customer Number::	22428	
---	-------	--

Domestic Priority Information

Application::	Continuity Type::	Parent Application::	Parent Filing Date::

Foreign Priority Information

Country::	Application number::	Filing Date::	Priority Claimed::

Assignee Information

Assignee name:: EnGeneIC Therapy Pty. Ltd.